

On Properties of Symmetric Banach Function spaces and Peetre's Interpolation Spaces

Eugene Tokarev

Dedicated to the memory of S. Banach.

ABSTRACT. A set $\mathcal{S}$ of all symmetric Banach function spaces defined on $[0,1]$ is equipped with a partial order by the relation $\subset^c$ of continuous inclusion. Properties of symmetric spaces, which does not depend of their position in the ordered structure are studied. With the help of the J. Peetre's interpolation scheme it is shown that for any pair E, F of symmetric spaces such that F is absolutely continuously included in E there exists an intermediate (so called, Peetre's) space $K(E, F; W)$, where W is a space with an unconditional basis, that is reflexive or, respectively, weakly sequentially complete if W also is reflexive or, respectively, weakly sequentially complete.

1. Introduction

A Banach space E of (classes of) Lebesgue measurable real functions acting on the interval $[0, 1]$ (equipped with the Lebesgue measure $\text{mes}(e)$ for a measurable subset $e \subset [0, 1]$) is called *symmetric* if for any pair of (measurable) functions $x = x(t)$ and $y = y(t)$ following conditions are satisfied:

1. If $x \in E$ and $|y(t)| \leq |x(t)|$ (a.e) then $y \in E$ and $\|y\|_E \leq \|x\|_E$.
2. If $x \in E$ and $|y(t)|$ is equimeasurable with $|x(t)|$ then $y \in E$ and $\|y\|_E = \|x\|_E$.

Let $\mathcal{S}$ be a class of all *symmetric Banach function spaces* (which in the future will be referred to as *symmetric spaces*). The class $\mathcal{S}$ may be partially ordered by a relation $E \subset^c F$, which means that E , as a set of functions, is contained in F and a norm $\|x\|_E$ of any function $x \in E$ is estimated by $\|x\|_E \leq c(E, F) \|x\|_F$. To be more precise, define an operator $i(E, F) : E \rightarrow F$, which asserts to any $x \in E$ the same function $x \in F$. This operator is called *the inclusion operator*. Its norm (which is equal to the infimum of all possible constants $c(E, F)$) is called *the inclusion constant*.

A lot of properties of symmetric spaces depend on their position in a partially ordered set $\langle \mathcal{S}, \subset^c \rangle$. E.g., extreme spaces, i.e., the minimal and the maximal elements of $\langle \mathcal{S}, \subset^c \rangle$, are exactly classical spaces $L_\infty[0, 1]$ and $L_1[0, 1]$ respectively (cf.

1991 *Mathematics Subject Classification*. Primary 46E30, 46B70; Secondary 46B10, 46B20.

Key words and phrases. Symmetric Banach spaces of measurable functions, Interpolation spaces, Partial order, Lattices.

[1]). Let G denotes an Orlicz space, which norming function is given by the equality $M(u) = (\exp(u^2) - 1)/(e - 1)$, and G_0 denotes a closure of $L_\infty[0, 1]$ in G . Any symmetric space E such that $G_0 \subset^c E \subset^c (G_0)^*$ has a basis of functions, which are uniformly bounded in the L_∞ -norm (cf. [2]). A lot of other similar examples may be found in [3], [4], [5].

Nevertheless there are properties of symmetric spaces E , which do not depend on position of E in the structure $\langle \mathcal{S}, \subset^c \rangle$. Some of such properties are studied in the article. In the study it will be used an interpolation construction due to J. Peetre [6], which asserts to a pair (E, F) of symmetric spaces with $F \subset^c E$ a new Banach space $K(E, F; W)$ (its construction uses a given Banach space W with an unconditional basis) that has certain interpolation properties with respect to spaces E and F .

This construction (and its generalizations) plays an important role in the functional analysis: in the interpolation theory (cf. [7]), in factorizations of weakly compact operators (cf. [8]), in constructing of peculiar examples of Banach spaces ([9], [10]) and many others.

The construction of the space $K(E, F; W)$ may one led to believe that its properties hardly depend on properties of E and of F .

However, if an order interval $[F, E]_{\subset^c}$ is *sufficiently large* (exactly: if F is *absolutely continuously included in* E ; shortly: $F \Subset E$; this relation will be explained later) then some properties of $K(E, F; W)$ depend only on properties of W . These properties will be called ω -properties and will be defined below. In particular, such is a property of a symmetric space to be either reflexive or weakly sequentially complete.

2. Definitions and notations

Let E be a symmetric space. Its *fundamental function* $\varphi_E(\tau)$ is given by

$$\varphi_E(\tau) = \|\chi_{[0, \tau]}(t)\|_E,$$

where $\chi_e(t)$ is an *indicator function* of a measurable subset $e \subset [0, 1]$:

$$t \in e \Rightarrow \chi_e(t) = 1; t \notin e \Rightarrow \chi_e(t) = 0$$

In the future it will be assumed that $\varphi_E(0) = 0$; $\varphi_E(1) = 1$.

E has an *absolutely continuous norm* provided

$$\lim_{\text{mes}(e) \rightarrow 0} (\|x(t)\chi_e(t)\|_E) = 0$$

for any $x(t) \in E$. It is known (see e.g. [11]) that E has such a norm if and only if it is separable.

A sequence $\{f_n : n < \infty\}$ may be also referred to as $(f_n)_{n < \infty}$ or, simply, as (f_n) . A sequence $(f_n) \subset E$ is said to be *disjoint* if $f_m(t)f_n(t) = 0$ (a.e.) for $n \neq m$.

Let E be a symmetric space, $x(t) \in E$, $\|x\|_E \neq 0$, $B \subset E$. Define following characteristics:

$$\eta_E(x, \tau) = \sup\{\|x(t)\chi_e(t)\|_E / \|x\|_E : \text{mes}(e) = \tau\};$$

$$\eta_E(B, \tau) = \sup\{\eta_E(x, \tau) : x \in B; \|x\|_E \neq 0\};$$

$$\eta_E(B) = \lim_{\tau \rightarrow 0+} \eta_E(B, \tau).$$

In a case when $E = L_1[0, 1]$ the lower index E will be omitted. Thus, $\eta(x, \tau)$ denotes $\eta_{L_1}(x, \tau)$ and so on.

Let E, F be a pair of symmetric spaces; $F \subset^c E$; $i : F \rightarrow E$ be the *inclusion operator*. It will be said that F is *absolutely continuously included in E* , in symbols: $F \Subset E$, if $\eta_E(iF) = 0$.

Let a, b be positive real numbers; E, F be symmetric spaces; f be a measurable function. Put

$$k(f; a, b) = \inf \{a \|u\|_E + b \|v\|_F : u + v = f\}.$$

Clearly, $k(f; a, b)$ may be regarded as a norm on a set $E + F$ of all functions f of kind $f = u + v$, for which $k(f; a, b)$ is finite.

DEFINITION 1. Let E, F be symmetric spaces; $F \Subset E$; W be a Banach space with an unconditional basis (w_i) . Let (a_i) and (b_i) be sequences of positive real numbers such that (a_i) is non decreasing; (b_i) is non increasing; $\lim_{i \rightarrow \infty} a_i = \infty$; $\sum_{i < \infty} b_i < \infty$.

A Peetre's space $K(E, F; W; (a_i), (b_i))$ is a Banach space of measurable functions $f(t)$, defined on $[0, 1]$, such that a series $\sum_{i < \infty} k(f; a_i, b_i) w_i$ convergents in W . A norm on the Peetre's space (that will be below referred to as $K(E, F; W)$ or, simply, as K) is given by

$$\|f\|_K = \left\| \sum_{i < \infty} k(f; a_i, b_i) w_i \right\|_W / \|\chi_{[0,1]}(t)\|_K.$$

Obviously, $K(E, F; W)$ is a symmetric space with an absolutely continuous norm (and, hence, is separable). As it was mentioned before, it will be supposed that $\varphi_K(1) = \|\chi_{[0,1]}(t)\|_K = 1$.

Let $(x_n) \subset X$ be a sequence of elements of a Banach space X . A sequence $(y_k) \subset X$ is said to be a *block sequence with respect to (x_n)* if there exist a sequence (a_n) of reals and a sequence $1 = n_0 < n_1 < \dots < n_k < \dots$ of naturals such that for all $k < \infty$

$$y_k = \sum_{m=n_k-1}^{n_{k+1}-1} a_m x_m$$

If (x_n) is a basic sequence then a block sequence (y_k) with respect to (x_n) will be called a *block-basis sequence*.

3. ω -properties of Peetre's spaces.

A property (P) of a Banach space X is said to be an ω -property if a condition "every subspace of X has a subspace that has the property (P)" implies that X also has the (P). The reflexivity (R) and the weak sequential completeness (WSC) are not ω -properties of general Banach spaces (take a look, e.g., at the classical James' quasireflexive space J). Nevertheless on the class $\mathcal{S}$ of symmetric spaces both of them, (R) and (WSC), are ω -properties. Indeed, if $E \in \mathcal{S}$ is not reflexive then it contains a subspace, that is isomorphic either to l_1 or to c_0 ; if E is not weakly sequentially complete then E contains a subspace which is isomorphic to c_0 . Clearly, every subspace of both l_1 and c_0 is non-reflexive; every subspace of c_0 fails to have the property (WSC).

THEOREM 1. Let E, F be symmetric spaces; $F \Subset E$; $(a_i); (b_i)$ be as in definition 1; W be a space with an unconditional basis (w_i) . Then every disjoint sequence of functions $(f_n) \subset K(E, F; W)$ contains a subsequence (f_{n_m}) , which is equivalent to a some block-basic sequence with respect to (w_n) .

PROOF. Let $x(t)$ be a measurable function. Put $s(\tau) = \eta_E(F, \tau)$. Since $F \in E$, it is clear that $\lim_{\tau \rightarrow 0+} s(\tau) = 0$. Let

$$A_n = \sum_{k=1}^n a_k.$$

Since $a_k \uparrow \infty$, for every $\varepsilon_0 > 0$ and every $n_0 \in \mathbb{N}$ there exists a number $j = j(\varepsilon_0, n_0)$ such that $2A_n a_j^{-1} < \varepsilon_0$. Fix n_0 and ε_0 and consider a function $f(t) \in K$; $\|f\|_K = 1$ with a support $A_f = \text{supp}(f) := \{t \in [0, 1] : f(t) \neq 0\}$. It will be assumed that the support of f is sufficiently small, namely: $s(\text{mes}(A_f)) \leq a_j^{-1} b_j$, where $j = j(\varepsilon_0, n_0)$ have been chosen before. It will be shown that a norm of the function f can be calculated up to ε_0 only by using vectors w_n with $n > n_0$, and that n_0 tends to infinity provided $\text{mes}(\text{supp}(f))$ tends to zero (this procedure will be called 'a cutting of the head'). Let $\|f\|_K = 1$. Then

$$\inf \{a_j \|u\|_E + b_j \|v\|_F : u + v = f\} \leq 1.$$

Thus, $\|u\|_E \leq a_j^{-1}$; $\|v\|_F \leq b_j^{-1}$.

Without loss of generality it may be assumed that

$$\text{supp}(u) \subset \text{supp}(f); \text{supp}(v) \subset \text{supp}(f).$$

Indeed, in a contrary case from the equality $u\chi_{A_f} + v\chi_{A_f} = f\chi_{A_f} = f$ and because of the symmetry, it follows:

$$\|u\chi_{A_f}\|_E \leq \|u\|_E; \quad \|v\chi_{A_f}\|_F \leq \|v\|_F.$$

Clearly, $\|f - v\|_E = \|u\|_E \leq a_j^{-1}$. From the triangle inequality it follows that

$$\begin{aligned} \|f\|_E &\leq \|u\|_E + \|v\|_E \leq a_j^{-1} + \|v\|_E \\ &\leq a_j^{-1} + b_j^{-1} \eta_K(v, \text{mes}(A_f)) \leq a_j^{-1} + b_j^{-1} a_j^{-1} b_j \leq 2a_j^{-1}. \end{aligned}$$

Since $F \in E$,

$$\max(a_n, b_n) k(f; 1, 1) \geq k(f; a_n, b_n).$$

Hence,

$$a_m^{-1} k(f; a_m, b_m) \leq \|f\|_E \leq 2a_j^{-1}.$$

Consequently,

$$k(f; a_m, b_m) \leq 2a_j^{-1} a_m.$$

With no loss of generality it may be assumed that the basis (w_n) is 1-unconditional, i.e. that $\|\sum a_i w_i\| \leq \sum |a_i|$ for any choice of scalars (a_i) . So,

$$\begin{aligned} \left\| \sum_{m=1}^n k(f; a_m, b_m) w_m \right\|_W &\leq \left\| \sum_{m=1}^n 2a_j^{-1} a_m w_m \right\|_W \\ &\leq 2 \left(\sum_{m=1}^n a_m \right) a_j^{-1} = 2A_n a_j^{-1} \leq \varepsilon_0. \end{aligned}$$

The series $\sum_{m < \infty} k(f; a_m, b_m) w_m$ convergents in W . Hence, there exists $N = N(\varepsilon_0)$ such that

$$\left\| \sum_{N < m < \infty} k(f; a_m, b_m) w_m \right\|_W \leq \varepsilon_0.$$

('a cutting of the tail'). Therefore,

$$\left\| \sum_{m < n} k(f; a_m, b_m) w_m + \sum_{N(\varepsilon) < m} k(f; a_m, b_m) w_m \right\|_W \leq 2\varepsilon_0.$$

So, up to $2\varepsilon_0$, the norm $\|f\|_K$ of a function f may be computed by using only the vectors $(w_i)_{i=n}^N$ provided its support is sufficiently small. Let $(f_n) \subset K$ be a disjoint sequence of functions. Clearly, $\lim_{n \rightarrow \infty} \text{mes}(\text{supp}(f_n)) = 0$.

Chose a subsequence $(f_{n_k}) \subset (f_n)$ by an induction.

Let $n_1 = 1$. Assume that $f_{n_1}, f_{n_2}, \dots, f_{n_{m-1}}$ are already chosen. Let n_k be the least number m such that $s(\text{mes}(\text{supp}(f_m))) \leq a_j^{-1}b_j$, where $j = j(m, 2^{-m}\varepsilon)$. Clearly, the sequence (f_{n_k}) spans a subspace of K , which is isomorphic to a subspace of W that is spanned by a some block sequence with respect to $\{w_i : i < \omega\}$. $\square$

COROLLARY 1. *Let $E, F, W, \{a_i; b_i : i < \omega\}$ be as in definition 1. If the space W is reflexive then $K(E, F; W)$ is reflexive too. If W is weakly sequentially complete then $K(E, F; W)$ also has this property.*

PROOF. If K is not reflexive then it contains a subspace Y , which is isomorphic either to l_1 or to c_0 . If K is not weakly sequentially complete then it contains a subspace Y , which is isomorphic to c_0 . In both cases K contains a disjoint sequence of functions that spanned a subspace isomorphic either to l_1 or to c_0 . However this contradicts with properties of W . $\square$

4. References

1. Semyonov, E.M. *Inclusion theorems for Banach spaces of measurable functions*, Dokl. AN SSSR, **156** (1964) 1294-1295 (in Russian)
2. Plichko, A.N. and Tokarev, E.V. *Bases of symmetric function spaces*, Math. Notes, **42**, no. 1/2 (1987) 630-634 (transl. from Russian)
3. Rodin, V.A. and Semyonov, E.M. *Rademacher series in symmetric spaces*, Analysis Math., **1:4** (1975) 207-222
4. Tokarev, E.V. *Subspaces of symmetric spaces of functions*, Funct. Anal. and Appl., **13** (1979) 152-153 (transl. from Russian)
5. Tokarev, E.V. *Complemented subspaces of symmetric function spaces*, Math. Notes, **32** (1982) 926-928 (transl. from Russian)
6. Peetre, J. *A new approach to interpolation spaces*, Studia Math., **34** (1970) 23-43
7. Beauzamy, B. *Espaces interpolationnelles réels; topologie et géométrie*, Lect. Notes in Math. **666** (1978) 1-90
8. Davis, W.T., Figiel, T., Johnson, W.B. and Pełczyński, A. *Factoring weakly compact operators*, Journ. Funct. Anal., **17** (1974) 317-327
9. Johnson, W.B., Maurey, B., Schechtman, G. and Tzafriri, L. *Symmetric structures in Banach spaces*, Memoirs Amer. Math. Soc., **217** (1979) 1-298

B.E. UKRECOLAN, 33-81 ISKRINSKAYA STR., 61005, KHARKIV-5, UKRAINE
 E-mail address: evtokarev@univer.kharkov.ua